



Original Article

Valine Consumption and Endurance Exercise Modified the Inflammation in Rat Model of Anxiety/Depression.

Mohammad-Hossein Beigi^{1}*

¹Silicon Hall: Micro/Nano Manufacturing Facility, Faculty of Engineering and Applied Science, Ontario Tech University, Ontario, Canada.

Submission date: 26-08-2022

Acceptance date: 13-02-2023

Abstract

Background: Depression and anxiety are disabling conditions with side effects on cognition, quality of life, memory, and mood. Anxiety is a dysregulated mood condition that could disturb cognition. The present study aimed to evaluate the effect of dietary interventions and endurance training in managing anxiety and depressive symptoms.

Methods: In this study, 30 rats were divided into five groups: Normal group, anxiety/depression group, rat model of anxiety/depression who consumed valine 3.0 $\mu\text{mol}/500 \mu\text{l}$ intraperitoneal injection once/day for eight weeks, rat model of anxiety/depression who conducted endurance training (32 m/min for 45 minutes/session, six days a week for eight weeks), and rat model of anxiety/depression who conducted endurance training (75% $\text{VO}_{2\text{max}}$) and consumed valine (3.0 $\mu\text{mol}/500 \mu\text{l}$). The level of anxiety/depression was evaluated by behavior open field tests and relative expression assay of the genes related to the inflammation.

Results: Findings indicated that the expression level of the NF- κB and TNF α significantly increased in the anxiety/depression group. Moreover, the endurance training and consumed valine improved the NF- κB and TNF α expression in the brain of the rat model of anxiety/depression. In addition, endurance training and consuming valine decreased the anxiety/depression level.

Conclusion: Endurance training and consuming valine reduced the brain's inflammation level and improved the anxiety/depression level.

Keywords: Anxiety, Depression Valine , Endurance training, Brain

* **Correspond Author:** mh_beigi10275@yahoo.com



Introduction

Psychiatric problems in both industrialized and developing countries have had a significant economic and social impact on healthcare systems (1, 2). The coronavirus pandemic could boost the risk of anxiety and depression because depression is a common mood disorder, which could lead to a rise in cerebrovascular disorders, heart diseases, gastrointestinal diseases, and the disruption of microbiota in the colon (3-5). The pathogenesis of depression is still unknown, but an artificial intelligence survey found some signaling pathways, including neuroinflammation, neurotransmitter function, phosphatidylinositol signaling system, gap junction, muscle contraction, insulin signaling pathway, and dopaminergic synapse, may be involved (5, 6).

The inflammation pathway could modulate behavior, the mucosal immune system, and emotion (7). Hence, managing or halting the inflammation might be essential for healthy brain function (8). Based on the evidence, depression and anxiety might be induced by cytokine storm and cytokine release syndrome (9). Furthermore, comprehensive studies have indicated that anxiety and depression could distribute sensory/motor neurons and increase brain inflammation (10). However, the mechanistic and molecular pathway is not elucidated.

The hippocampal area in depression is less than in healthy persons, according to Darren W.Roddy *et al.* Furthermore, they have found a link between the severity of depression and the size of the hippocampus (11). Hence, the hippocampus might play a decisive role in anxiety and depression.

No effective therapy strategy is available to stop the progression of depression. Of the therapies for depression, only 60% showed improvement in symptoms (12). As a result, an effective and safe treatment could shed light on how to reduce this illness and improve the gastrointestinal environment. In addition, epidemiological research suggests that environmental factors may play a significant role in the development of depression (13). Thus, the present study aimed to evaluate the effect of dietary interventions and endurance training in managing anxiety and depressive symptoms (14).

In addition, it was well-known that the production of neurotransmitters could be hampered without adequate physical activity and nutrition (15). Growing evidence demonstrated that regular exercise could improve inflammation agents and cytokine release syndrome (16, 17). Non-pharmacological interventions such as exercise effectively treat anxiety and depression (18). Mental disorders might be reduced by increased physical

exercise, probably due to their favorable impacts on psychological well-being, coping mechanisms, and self-perceptions of one's physical appearance (19).

An exciting aspect of depression treatment is that producing neurotransmitters by aromatic amino acids like phenylalanine, tyrosine, and tryptophan might be practical (20, 21). Evidence suggests that the blood-brain barrier (BBB) competes with valine for the transport of aromatic amino acids, which could lead to reduced neurotransmitter concentrations in aromatic amino acids and a delay in serotonin production, as well as a suppression of central tiredness. It is not apparent how valine amino acid concentrations affect depression, even though this is a common finding (22).

The study aimed to evaluate the significant hub genes in brain rats with depression-like behaviors. Moreover, we revealed that the hub genes involved in inflammation could improve with endurance training and Valine /Isovaline supplementation.

Materials and Methods

Animal Study

Male rats aged seven weeks were furnished and maintained in the animal facility. Rats were kept under conventional conditions of $24\pm 3^{\circ}\text{C}$ temperature, 55 to 60% humidity, 12-hour light and dark cycles, and ad libitum access to food and drink. In addition, all animals were provided with a regular diet. Before beginning the research, rats were acclimated for one week. In this study, the rats were distributed into five groups: Normal group, anxiety/depression group (An/Dep), rat model of anxiety/depression who consumed valine (Val) ($3.0\ \mu\text{mol}/500\ \mu\text{l}$ intraperitoneal injection once a day for eight weeks), rat model of anxiety/depression who conducted endurance training (Exercise) (32 m/min for 45 minutes each session, six days a week for eight weeks), and rat model of anxiety/depression who conducted endurance training and consumed valine (Ex+Val). The infusion of xylazine (10 mg/kg body weight per animal) and ketamine (80 mg/kg body weight per mouse) were used to euthanize rats.

Inducing Depression

As stated previously, we separated rats with depression into four groups. Rats were treated with fixation, electric tail, and foot shocks for 1 second with ten repetitions (0.5 mA intensity) and food deprivation, followed by 14 days. (23, 24).

Rat behavioral tests

Open field test

Rest periods and locomotor activity were assessed. The open field space was split into 16 squares measuring 40×40×40 cm. Each mouse was positioned within the core area and observed for one-hour periods. Distance moved (cm), duration time (sec), and Rest times (sec) were computed and scored (25, 26).

Valine complement and food intake

Rats had unrestricted access to regular food and running water. In addition, 3.0 μmol/500 μl of valine was administered intraperitoneally once every day for two months (25, 27).

Exercise protocol

On a motorized treadmill, endurance training (EX) was executed. On the treadmill, the intensity was moderate to high for two months (6 days a week). After that, the workout and running pace length gradually increased to 75% VO₂ max (32 m/min) for 45 minutes. Additionally, the slope of the treadmill was deemed to be 0% (28, 29).

Quantitative real-time PCR (qRT-PCR)

The TRIzol reagent isolated total RNA from the hippocampal area and ileum tissue (Sigma, USA). cDNA synthesis was performed using 1 g of total RNA and a cDNA synthesis kit according to the manufacturer's instructions (TaKaRa, Japan). qRT-PCR was performed with CYBR Green (TaKaRa, Japan) and Corbet rotor gene 6000 (Qiagen, Australia). The detection of gene expression was assessed using the $2^{-\Delta\Delta CT}$ technique. The expression levels of glyceraldehyde-3-phosphate dehydrogenase (Gapdh) were used to compute the relative expression of genes.

Statistical Analysis

Using GraphPad Prism, statistical analysis was performed (Version 9; GraphPad Software). Distribution was normalized using the Shapiro-Wilk test, and variables were normally distributed. Due to numerous comparisons, the data were analyzed using a one-

way analysis of variance (ANOVA) with Tukey's post hoc test. At $P.value < 0.05$, differences were judged statistically significant. In addition, data were presented as the mean \pm SD.

Result

This study indicated that the total movement rat model of anxiety/depression was reduced compared with the Normal group (Figure 1). Moreover, the endurance training increased the total movement compared with the An/Dep group. In addition, consuming valine enhanced the total movement compared with the An/Dep group. Furthermore, the endurance training and consuming valine (Ex+Val) significantly elevated the total movement compared with the other group.




Figure 1. Estimated total distance movement (Unit) via open field test. (mean \pm SD; n = 6)

The results revealed that the expression level of the NF- κ B and Tnf α in the anxiety/depression group was amplified compared with the control group in the hippocampus (Figure 2). Also, endurance training and valine consumption diminished the relative expression of the level of the NF- κ B and Tnf α in the hippocampus (Figure 2). On the other hand, endurance training along with valine consumption decreased the expression level of the NF- κ B and Tnf α compared with the other groups (Figure 2).




Figure 2. Expression level of the Tnf α and NF- κ B. (mean \pm SD; n = 6)

Discussion

In this study, we indicated that the inflammation agents such as NF- κ B and Tnf α significantly increased in the rat model of anxiety/depression hippocampus. Moreover, we found that endurance training along with valine consumption regulated the expression level of the NF- κ B and Tnf α compared with the other groups. Literature review showed that several molecular mechanisms, such as Toll-Like receptor, Cytokine-Cytokine, IL-17 pathway, and TNF signaling pathway involved in the pathomechanism of anxiety/depression (30). In addition, emerging data suggested that receptors of the advanced glycation end product are elevated in neurodegenerative diseases compared to samples from normal patients (31). These results from earlier investigations confirmed that inflammation is a neuropathologic change facilitator (31). A separate investigation determined that increased inflammation such as NF- κ B and Tnf α might induce depression condition [56]. These findings showed that the Toll-Like receptor, Cytokine-Cytokine, IL-17 pathway, and TNF signaling pathway may play a crucial role in anxiety/depression. These signalings have been recognized as one of the prospective treatment methods for depression (32). Since proliferation-differentiation and cell cycle programming are essential for body homeostasis, Cytokine-Cytokine and TNF signaling pathways may be useful in determining the cell destiny of many organs (33). Therefore, Cytokine-Cytokine and TNF signaling pathways were altered in the hippocampus regions of mice with depression. In the present study, we demonstrated that the expression level of the mRNAs involved the inflammation signaling

pathway. Based on our data, we found that the expression level of the NF- κ B and Tnf α significantly increased in the depression condition. Also, we established that receiving valine could improve the expression of the NF- κ B and Tnf α in the depression condition. Several studies have indicated that consumption of the valine has an advantageous effect and can impact molecular processes. This natural supplement can be alternative and complementary medicine in preventing and treating the symptoms of a disease in conjunction with a prescription drug. Based on these findings, valine may be a safe and useful adjunct in treating several neurological and psychiatric illnesses. Evidence has demonstrated that valine, leucine, and isoleucine consumption altered the involved metabolism and mitochondrial biogenesis gene expression (29).

Moreover, Deyang Yu *et al.* revealed valine could mediate metabolic health and regulate the FGF21. Based on these results, branched-chain amino acids could improve energy expenditure and induce Ucp1 (27). Notably, our study revealed that endurance exercise improved the expression of NF- κ B and Tnf α . Immense evidence has indicated that endurance exercise is considered non-pharmacological interference. Based on the epidemiology studies, endurance exercise might regulate inflammation, antioxidant balance, subtractive lipid-peroxidation, and oxidative stress status (16, 17). Although the particular pathomechanism involved in inflammation has not been fully understood, artificial intelligence analysis and text mining emphasized that high oxidative stress, apoptosis, and inflammation increased depression risk (18). Paolucci and colleagues indicated that moderated intensity interval training decreased the concentration of the Tnf α and interleukin-1 beta (IL-1 β), C-reactive protein (CRP), and interleukin-6 (IL-6). Moreover, they found that 6 weeks of moderated intensity interval training modified the student's mental health (34). In another study, exercise significantly decreased the concentration of the interleukin-1 β (IL-1 β), interleukin-6 (IL-6), Tnf α , and interferon- γ (IFN- γ) of patients involved in the major depressive disorder. Based on these data, physical activity could have the antidepressant medications (35)

Conclusion

According to biological analysis approaches, we could conclude that synchronizing valine consumption and endurance exercise could have a synergetic effect on the

inflammation level. Hence, endurance training and consuming valine reduced the brain's inflammation level and improved the anxiety/depression level.

Competing interests

There is no competing of interest to disclose.

Acknowledgements

We thank our colleagues for their association and helpful discussions in this study.

References:

- Jadhakhan F, Lindner OC, Blakemore A, Guthrie E. Prevalence of common mental health disorders in adults who are high or costly users of healthcare services: protocol for a systematic review and meta-analysis. *BMJ open*. 2019;9(9):e028295.
- Krishnan V, Nestler EJ. The molecular neurobiology of depression. *Nature*. 2008;455(7215):894-902.
- Hawes MT, Szenczy AK, Klein DN, Hajcak G, Nelson BD. Increases in depression and anxiety symptoms in adolescents and young adults during the COVID-19 pandemic. *Psychological Medicine*. 2021:1-9.
- Indriani S, Trisyani Y, Lumbantobing VB. DEPRESSION IN PATIENT WITH HEART FAILURE: A SCOPING REVIEW. *UNEJ e-Proceeding*. 2021:11-28.
- Bailey MT. Influence of stressor-induced nervous system activation on the intestinal microbiota and the importance for immunomodulation. *Microbial endocrinology: the microbiota-gut-brain axis in health and disease*. 2014:255-76.
- Carabotti M, Scirocco A, Maselli MA, Severi C. The gut-brain axis: interactions between enteric microbiota, central and enteric nervous systems. *Annals of gastroenterology: quarterly publication of the Hellenic Society of Gastroenterology*. 2015;28(2):203.
- Aoki-Yoshida A, Aoki R, Moriya N, Goto T, Kubota Y, Toyoda A, et al. Omics studies of the murine intestinal ecosystem exposed to subchronic and mild social defeat stress. *Journal of proteome research*. 2016;15(9):3126-38.
- Borre YE, O'Keeffe GW, Clarke G, Stanton C, Dinan TG, Cryan JF. Microbiota and neurodevelopmental windows: implications for brain disorders. *Trends in molecular medicine*. 2014;20(9):509-18.
- Abedpoor N, Taghian F, Hajibabaie F. Physical activity ameliorates the function of organs via adipose tissue in metabolic diseases. *Acta Histochem*. 2022;124(2):151844.
- Foster J, Neufeld K-A, editors. *Gut-brain axis: how the microbiome influences anxiety and depression*. International Journal of Neuropsychopharmacology; 2014: CAMBRIDGE UNIV PRESS 32 AVENUE OF THE AMERICAS, NEW YORK, NY 10013-2473 USA.
- Roddy DW, Farrell C, Doolin K, Roman E, Tozzi L, Frodl T, et al. The hippocampus in depression: more than the sum of its parts? Advanced hippocampal substructure segmentation in depression. *Biological Psychiatry*. 2019;85(6):487-97.

- Wang PS, Lane M, Olfson M, Pincus HA, Wells KB, Kessler RC. Twelve-month use of mental health services in the United States: results from the National Comorbidity Survey Replication. *Archives of general psychiatry*. 2005;62(6):629-40.
- Network T, Consortium PASotPG. Psychiatric genome-wide association study analyses implicate neuronal, immune and histone pathways. *Nature neuroscience*. 2015;18(2):199.
- Firth J, Marx W, Dash S, Carney R, Teasdale SB, Solmi M, et al. The effects of dietary improvement on symptoms of depression and anxiety: a meta-analysis of randomized controlled trials. *Psychosomatic medicine*. 2019;81(3):265.
- Maher TJ. Effects of nutrients on brain function. *Progress in brain research*. 2000;122:187-94.
- Adeniyi AF, Okafor NC, Adeniyi CY. Depression and physical activity in a sample of nigerian adolescents: levels, relationships and predictors. *Child and adolescent psychiatry and mental health*. 2011;5(1):1-10.
- Taliaferro LA, Rienzo BA, Pigg RM, Miller MD, Dodd VJ. Associations between physical activity and reduced rates of hopelessness, depression, and suicidal behavior among college students. *Journal of American College Health*. 2009;57(4):427-36.
- Mammen G, Faulkner G. Physical activity and the prevention of depression: a systematic review of prospective studies. *American journal of preventive medicine*. 2013;45(5):649-57.
- Knapen J, Vancampfort D, Moriën Y, Marchal Y. Exercise therapy improves both mental and physical health in patients with major depression. *Disability and rehabilitation*. 2015;37(16):1490-5.
- McLean A, Rubinsztein JS, Robbins TW, Sahakian BJ. The effects of tyrosine depletion in normal healthy volunteers: implications for unipolar depression. *Psychopharmacology*. 2004;171(3):286-97.
- Firk C, Markus CR. Serotonin by stress interaction: a susceptibility factor for the development of depression? *Journal of Psychopharmacology*. 2007;21(5):538-44.
- Fernstrom JD. Branched-chain amino acids and brain function. *The Journal of nutrition*. 2005;135(6):1539S-46S.
- Wang Q, Timberlake II MA, Prall K, Dwivedi Y. The recent progress in animal models of depression. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2017;77:99-109.
- Tian P, O'Riordan KJ, Lee Y-k, Wang G, Zhao J, Zhang H, et al. Towards a psychobiotic therapy for depression: *Bifidobacterium breve* CCFM1025 reverses chronic stress-induced depressive symptoms and gut microbial abnormalities in mice. *Neurobiology of stress*. 2020;12:100216.
- Goto T, Kubota Y, Tanaka Y, Iio W, Moriya N, Toyoda A. Subchronic and mild social defeat stress accelerates food intake and body weight gain with polydipsia-like features in mice. *Behavioural brain research*. 2014;270:339-48.
- Kim YR, Park B-K, Kim YH, Shim I, Kang I-C, Lee MY. Antidepressant effect of *fraxinus rhynchophylla* hance extract in a mouse model of chronic stress-induced depression. *BioMed research international*. 2018;2018.
- Yu D, Richardson NE, Green CL, Spicer AB, Murphy ME, Flores V, et al. The adverse metabolic effects of branched-chain amino acids are mediated by isoleucine and valine. *Cell metabolism*. 2021;33(5):905-22. e6.

- Schefer V, Talan MI. Oxygen consumption in adult and AGED C57BL/6J mice during acute treadmill exercise of different intensity. *Experimental gerontology*. 1996;31(3):387-92.
- Abedpoor N, Taghian F, Ghaedi K, Niktab I, Safaiejad Z, Rabiee F, et al. PPAR γ /Pgc-1 α -Fndc5 pathway up-regulation in gastrocnemius and heart muscle of exercised, branched chain amino acid diet fed mice. *Nutrition & metabolism*. 2018;15(1):1-15.
- Li Z, Pham T, Tamir H, Chen J, Gershon M. Enteric dopaminergic neurons: definition, developmental lineage, and effects of extrinsic denervation. *Journal of Neuroscience*. 2004;24(6):1330-9.
- Li Y, Peng Y, Shen Y, Zhang Y, Liu L, Yang X. Dietary polyphenols: regulate the advanced glycation end products-RAGE axis and the microbiota-gut-brain axis to prevent neurodegenerative diseases. *Critical Reviews in Food Science and Nutrition*. 2022:1-27.
- Singhal G, Jawahar MC, Morgan J, Corrigan F, Jaehne EJ, Toben C, et al. TNF signaling via TNF receptors does not mediate the effects of short-term exercise on cognition, anxiety and depressive-like behaviors in middle-aged mice. *Behavioural Brain Research*. 2021;408:113269.
- Liu H, Huang X, Li Y, Xi K, Han Y, Mao H, et al. TNF signaling pathway-mediated microglial activation in the PFC underlies acute paradoxical sleep deprivation-induced anxiety-like behaviors in mice. *Brain, Behavior, and Immunity*. 2022;100:254-66.
- Paolucci EM, Loukov D, Bowdish DM, Heisz JJ. Exercise reduces depression and inflammation but intensity matters. *Biological psychology*. 2018;133:79-84.
- Rethorst CD, Toups MS, Greer TL, Nakonezny PA, Carmody TJ, Grannemann BD, et al. Pro-inflammatory cytokines as predictors of antidepressant effects of exercise in major depressive disorder. *Molecular psychiatry*. 2013;18(10):1119-24.