

# Influence of *Spirulina* sp. and citric acid dietary supplements on the growth performance and immune parameters of common carp (*Cyprinus carpio*)

Hossein Sabetmand . Hamid Faghani Langarudi  . Abbasali Zamini . Babak Tizkar

Received: 17 November 2022 / Accepted: 14 April 2023 / Published online: 22 February 2024  
© The Author(s) 2024

**Abstract** This study explores the effects of supplementing the diets of common carp (*Cyprinus carpio*) with *Spirulina* sp. and citric acid (CA) on growth performance and immune parameters. Various experimental diets were formulated, including a control group, diets with 20g/kg *Spirulina*, 30g/kg *Spirulina*, 0.5g/kg citric acid, 1.0g/kg citric acid, and four mixed combinations of *Spirulina* and citric acid. Initially, common carp weighing  $15.3 \pm 1.9$ g were fed these diets for eight weeks in 40-L aquaria with three replicates. Growth performance and immune indices, specifically lysozyme, IgM, and immunoglobulin M (IgM) levels were assessed. The results demonstrated that the *Spirulina*/citric acid mixture significantly improved growth performance and immune indices compared to individual *Spirulina* and citric acid treatments and the control group ( $P < 0.05$ ). The growth indices indicated increased dietary efficiency with the higher inclusion of the mixture, with the 30g *Spirulina* sp. + 1.0g citric acid and 30g *Spirulina* sp. + 0.5g citric acid treatments exhibiting the highest growth performance and the lowest feed conversion ratio (FCR). In conclusion, dietary supplementation of 30g/kg *Spirulina* combined with 0.5g/kg citric acid was found to promote growth and positively influence the immune parameters in common carp.

**Keywords** *Spirulina* sp. . *Cyprinus carpio* . Growth . Immune . Digestive enzymes

## Introduction

Aquaculture has witnessed remarkable growth in animal protein production in recent decades. Among the most frequently cultivated fish worldwide is the common carp (*C. carpio*). This fish holds a significant position in the aquaculture industry of many Asian and some European countries. Notably, it is a valuable source of nutrients, making it a key component of healthy human diets. The common carp is highly prized for its numerous desirable traits, including rapid growth, efficient conversion of natural and supplementary feeds, and relative resilience to adverse environmental conditions and diseases. Proper nutrition is essential to ensure this cultivated fish's high survivability and accelerated growth rate.

Algae have recently emerged as pivotal food sources and additives in the commercial rearing of various aquatic animals, mainly fish and penaeid prawn larvae (Borowitzka 1997; Belay et al. 1996; Khatoun et

---

Hossein Sabetmand  
PhD Student of Fisheries Sciences, Department of Fisheries, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran

Hamid Faghani Langarudi (✉)  
Department of Fisheries, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran  
e-mail: hamid\_faghani1@yahoo.com

Abbasali Zamini  
Department of Fisheries, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran; Department of Fisheries, Lahijan Branch, Islamic Azad University, Lahijan, Iran

Babak Tizkar  
Department of Fisheries, Tonekabon Branch, Islamic Azad University, Tonekabon, Iran

al. 2010). Multiple studies have evaluated the nutritional value of dried microalgae as feed components for crustaceans and fish larvae (Biedenbach et al. 1990; Navarro and Sarasquete 1998; Khatoon et al. 2009). Kumar et al. (2010) analyzed the impact of periphyton and supplementary nutrition, using commercial pelleted feeds, on the growth performance of juvenile Nile tilapia *Oreochromis niloticus*. This lower plant group contains many vitamins, pigments, and nearly all essential nutrients, including polyunsaturated fatty acids (PUFA). It also serves as a valuable source of proteins and carbohydrates. Given the profound influence of various algae in general on the growth and vitality of fish, numerous algae species have been confirmed over time. Algae genera such as *Spirulina* sp. are widely embraced in aquaculture feeds due to their remarkable nutritional content (Avron and Ben-Amotz 1992; Lee 1997; Yamaguchi 1997).

*Spirulina*, in particular, has gained recognition as a valuable supplementary ingredient in aqua feed owing to its high protein and vitamin content. Microalgae typically boast a crude protein content of approximately 50%, with an amino acid profile akin to fishmeal. Furthermore, they serve as a source of polyunsaturated fatty acids (PUFAs), particularly those from the n-3 series. Consequently, microalgae have the potential to replace fish oil and fish meal in diets, enhancing meat quality through the deposition of n-3 PUFAs (Roy and Pal 2015; Sarker et al. 2016). Microalgae, known for their oil and molecule content, can also be harnessed as feed stocks for producing biofuels and high-value products, representing a promising renewable energy source (Moreno-Garcia et al. 2017). *Spirulina*, a commonly cultivated microalgae species in the commercial sector (Priyadarshani and Rath 2012; Farag et al. 2016), is a type of cyanobacteria renowned for its high protein content, ranging from 56% to 69% by dry weight. Additionally, *Spirulina* is replete with various bioactive compounds, including minerals, vitamins, carotenes, essential fatty acids, and antioxidants (Venkataraman 1997). Moreover, it serves as an effective substitute for animal-derived proteins in the diets of fish species, demonstrating favorable outcomes across a spectrum of species such as *Oreochromis niloticus* (Lu and Takeuchi 2004), *Paralichthys olivaceus* (Kim et al. 2015), *Clarias batrachus* (Dar et al. 2014), *Oplegnathus fasciatus* (Rahimnejad 2013), *Cirrhinus mrigala* (James et al. 2009), and *Poecilia reticulata* (Dernekbasi et al. 2010).

A significant portion of phosphate (P), roughly 60% to 70%, in vegetable protein ingredients is bound to citric acid. This binding may hinder the availability of P when there is an increase in dietary vegetable protein compounds and minerals like zinc, magnesium, and calcium (Denstadli et al. 2010). Adding citric acid (CA) to diets has enhanced P release from phytate in vitro (Zyla et al. 1995). Lowering the intestinal pH increases the solubility of P and phytate, thereby enhancing P absorption in the small intestine (Cross et al. 1990). Furthermore, apart from its impact on intestinal pH, the interaction between supplementary organic acids and various cations along the intestine can serve as a chelating agent, resulting in improved intestinal mineral absorption (Wood and Serfaty-Lacrosniere 1992). While extensive research exists regarding the effect of dietary acidification on mineral utilization in terrestrial animals, studies involving fish have been somewhat limited (Sarker et al. 2005). Experiments conducted with pigs have indicated that CA supplementation promotes growth performance (Sugiura et al. 2001). In the case of rainbow trout, citric acid supplementation in their diet has been observed to chelate CA and P, increasing their solubility and enhancing mineral utilization (Vielma et al. 1999).

Therefore, microalgae can be directly employed as a live culture or a value-added dietary supplement. It is imperative to recognize that dietary patterns significantly influence the development and functionality of the immune system. Hence, any newly formulated algae diet must meet the specific nutritional requirements of the fish. In the present investigation, *Spirulina* algae and citric acid were isolated, blended at varying levels, and assessed over 8 weeks to discern which combinations yielded superior growth rates and immune system performance.

## Materials and methods

### Fish and experimental procedure

This study was conducted at the Research and Education Center for Natural Resources and Agriculture at the Fisheries Campus in Guilan Province, Iran. The adaptation process for the fish involved housing them in a 2000-liter tank and providing them with the prescribed dietary regimen for one week. Following this acclimatization, the fish were relocated to twenty-seven 40-liter aquaria equipped with central aeration. To



ensure the water quality remained suitable for the fish, physicochemical parameters were monitored using a dial oximeter (OXI3230B/SET) and a pH meter (PH330i/SET), with measurements taken three times daily.

#### Experimental design, diets, and performance

A commercial carp feed supplied by Dan-Vahdat Co., Iran, was employed in the experiments. The feed consisted of 35% crude protein and included wheat flour, wheat bran, fish meal, soybean meal, corn, and rice bran. Additionally, the feed was supplemented with *Spirulina* and citric acid.

*Spirulina* powder and citric acid were evenly distributed onto the feed and dried at ambient temperature. Analysis of the fish diet revealed the following nutritional content: crude protein at 35%, fat at 11%, fiber at 10%, ash at 14%, calcium at 6%, phosphorus at 9%, and moisture at 14%. The ingredients included in the fish meal were corn meal, soybean meal, corn gluten, soy protein, concentrated wheat meal, soybean oil, fish meal, sodium chloride, choline chloride, vitamin and mineral premixes, probiotics, amino acids, and antioxidants.

The experimental methodology encompassed a control group and several treatment groups, each denoted by specific quantities of *Spirulina* powder and citric acid per kilogram of diet. These treatments were as follows: 0g of *Spirulina* powder and 0g of citric acid per kilogram of diet (SpCA-0), 20g of *Spirulina* (Sp20), 30g of *Spirulina* (Sp30), 0.5g of citric acid (CA0.5), 1g of citric acid (CA1), 20g of *Spirulina* + 0.5g of citric acid (Sp20+CA0.5), 20g of *Spirulina* + 1g of citric acid (Sp20+CA1), 30g of *Spirulina* + 0.5g of citric acid (Sp30+CA0.5), and 30g of *Spirulina* + 1g of citric acid (Sp30+CA1) per kilogram.

The fish received two daily feedings, and biometric measurements were taken at the beginning, middle, and end of the experimental period, with a 24-hour interval after feeding cessation. Randomly selected fish had blood samples collected and transferred to the hematology laboratory for immunological assessment. A comprehensive database was maintained, with monthly biometric measurements, including fish body length and weight data.

#### Measurement of growth performance

Statistical calculations on growth indices and food efficiency were conducted based on the following formulas:

BWI (*Body weight index*%) = 100% (Wt-W0) / W0 [W0 = Mean initial weight (g); Wt = Average final weight (g)]

FCR (Food conversion ratio) = F / (Wt-W0) [F = the amount of food consumed by fish; W<sub>0</sub> = Mean primary biomass (g); W<sub>t</sub> = Mean final biomass (g)]

SGR (Specific growth rate) = (ln Wt - ln W0) / t × 100 [W0 = Mean primary biomass (g); W<sub>t</sub> = Mean final biomass (g); T = Period (days); ln is the natural log, and t stands for the experimental period in days]

#### Immunological factors

The investigation of immunological factors was conducted during the final phase of this research, which took place in the eighth week. Blood samples from the fish were randomly collected for analysis. To ensure a stress-free process, the fish were initially anesthetized using clove powder, and their blood was then extracted from the caudal vasculature using insulin syringes. A total of ten blood specimens were obtained from each replicate group. The exact number of samples was collected from the treatment and control groups, each with three replicates. Subsequently, the blood samples were processed to obtain blood serum through centrifugation.

The Immunoglobulin M (IgM) levels were quantified using the immunoturbidimetric method. In this method, IgM forms a complex with polyclonal antibodies in buffer solutions, causing the solution to become turbid. The degree of turbidity is directly proportional to the concentration of IgM and was measured using a spectrophotometer, with distilled water serving as the blank (Model 2100-VIS by Unico USA) at a wavelength of 340 nm. The determination of serum lysozyme levels involved utilizing 1.75 ml of micrococcus lysodeikticus (Sigma) suspension, equivalent to 0.375 mg/ml sodium phosphate buffer at 0.05 M and pH 6.2. To this suspension, 250 microliters of mixed serum samples were added, and light absorption



readings were taken after 15 and 180 seconds using a spectrophotometer at a wavelength of 670 nm. A blank was established using sodium phosphate buffer. The quantification of total immunoglobulin concentrations was performed according to the methodology delineated by Siwicki and Anderson in 1993 and the approach outlined by Amar et al. in 2000. In brief, the serum sample underwent the Biuret method. More specifically, 0.1 ml of each serum sample was mixed with 0.1 ml of a 32% polyethylene glycol solution (PEG, 10,000 MW, Sigma Chemical, St. Louis, MO, USA). This resultant mixture was subsequently incubated for 2 hours, facilitating the precipitation of immunoglobulin molecules. The precipitated immunoglobulin was separated through centrifugation (Eppendorf Centrifuge 5415R, Eppendorf AG, Hamburg, Germany) at 5000 revolutions per minute and a temperature of 4°C. The supernatant fluid was subjected to measurement for total protein content, and the immunoglobulin concentration was computed employing the subsequent formula: Total Immunoglobulin Concentration (mg per ml) = Total protein treated with polyethylene glycol - Total protein in serum samples. This approach allowed us to determine the immunoglobulin concentration in milligrams per milliliter.

### Data analysis

The normality of the data was evaluated utilizing the Shapiro-Wilk test. To ascertain the significance of the experiments, a one-way ANOVA analysis was conducted employing the Duncan method at a 95% confidence level ( $P < 0.05$ ). The data analysis was executed using SPSS 20.0 and Excel 2010 software.

## Results

### Growth parameters analysis

The study outcomes reveal distinct daily growth rates among various treatment groups. Notably, the highest daily growth rate was observed in SP30+CA0.5 ( $0.13 \pm 0.06$ ) and SP30+CA1 ( $0.13 \pm 0.03$ ), while the lowest was associated with SP20 ( $0.08 \pm 0.01$ ). Statistical analysis demonstrates significant variance between the treatments throughout the experiment ( $P < 0.05$ ) (Table 1). Discriminant testing further indicates a significant divergence in the daily growth coefficient of SP30+CA0.5 compared to other treatments. However, it exhibited no statistical distinctions from SP20+CA0.5, SP20+CA1, and SP30+CA1 ( $P > 0.05$ ). Additionally, the results suggest that SP20, SP30, CA0.5, and CA1 share a statistically equivalent daily growth coefficient ( $P > 0.05$ ).

Findings show considerable differences in weight gain rates among the treatment groups. Specifically, the highest weight gain rate was observed in SP30+CA1 ( $8.7 \pm 0.31$ ), whereas the lowest was associated with SP20 ( $4.76 \pm 0.61$ ) (Table 1). Statistical assessments reveal significant differences in weight gain among the treatments during the experimental period ( $P < 0.05$ ). Discriminant testing indicates a noteworthy difference in the weight gain of SP30+CA1 compared to other treatments. Nonetheless, SP30+CA1 did not display a statistically significant difference from SP30, SP20+CA0.5, SP20+CA1, and SP30+CA0.5 ( $P > 0.05$ ). Furthermore, results indicate that SP20, CA0.5, and CA1 statistically share the same weight gain ( $P < 0.05$ ).

The study demonstrates variations in the percentage of body weight index among the treatment groups. The highest percentage of body weight index is associated with SP30+CA1 ( $35 \pm 91 \pm 1.47$ ), while the lowest is related to SP20 ( $21.22 \pm 2.98$ ) (Table 1). Statistical analysis indicates significant differences in body weight index among the treatments during the experimental period ( $P < 0.05$ ). SP30+CA1, however, does not exhibit any significant differences from SP20+CA1 and SP30+CA0.5 ( $P > 0.05$ ). Results further reveal that SP20, SP30, CA0.5, and CA1 statistically share the same percentage of body weight index ( $P > 0.05$ ).

The investigation unveils varying specific growth coefficients among the treatment groups. Specifically, the highest specific growth coefficient is associated with SP30+CA1 ( $0.178 \pm 5.11$ ), while the lowest is linked to SP20 ( $0.40 \pm 3.2$ ) (Table 1). Statistical tests demonstrate significant differences in specific growth rates among the treatments throughout the experiment ( $P < 0.05$ ). According to the findings, the specific growth coefficient of SP30+CA1 significantly differs from that of other treatments ( $P < 0.05$ ). However, no statistical differences are observed with SP20+CA0.5, SP20+CA1, and SP30+CA0.5 ( $P > 0.05$ ). Additionally, results indicate that SP20, SP30, CA0.5, and CA1 share the same specific growth rate (Table. 1).



**Table 1** Effect of Spirulina and CA supplementation on growth performance of common carp in different treatments  
Spirulina and CA concentrations g per kg (food)

	Spirulina						CA			Mixture (Spirulina+CA)		
	Control	20	30	0.5	1.0	20+0.5	20+1.0	20+1.0	30+0.5	30+1.0	30+1.0	
W <sub>0</sub> (g)	22.20±3.70 <sup>a</sup>	22.40±6.28 <sup>a</sup>	22.60±5.36 <sup>a</sup>	24.00±4.83 <sup>a</sup>	22.90±6.10 <sup>a</sup>	23.60±5.13 <sup>a</sup>	22.70±8.10 <sup>a</sup>	22.80±5.1 <sup>a</sup>	22.80±5.1 <sup>a</sup>	22.70±6.38 <sup>a</sup>		
W <sub>t</sub> (g)	24.00±8.34 <sup>a</sup>	25.60±7.72 <sup>a</sup>	26.38±4.57 <sup>a</sup>	23.80±3.61 <sup>a</sup>	26.25±2.95 <sup>a</sup>	25.10±4.95 <sup>a</sup>	27.50±11.34 <sup>a</sup>	25.44±6.39 <sup>a</sup>	25.44±6.39 <sup>a</sup>	26.44±5.43 <sup>a</sup>		
GR <sub>(g)</sub>	0.08±0.01 <sup>b</sup>	0.08±0.01 <sup>b</sup>	0.10±0.01 <sup>bc</sup>	0.09±0.01 <sup>cd</sup>	0.10±0.01 <sup>cd</sup>	0.12±0.01 <sup>ab</sup>	0.12±0.01 <sup>a</sup>	0.13±0.01 <sup>a</sup>	0.13±0.01 <sup>a</sup>	0.13±0.01 <sup>a</sup>		
W <sub>C</sub> (g)	4.82±0.59 <sup>b</sup>	4.76±1.05 <sup>b</sup>	6.08±0.78 <sup>bc</sup>	5.15±0.07 <sup>cd</sup>	5.79±0.52 <sup>cd</sup>	6.98±0.50 <sup>b</sup>	7.51±0.49 <sup>a</sup>	7.70±0.73 <sup>a</sup>	7.70±0.73 <sup>a</sup>	8.06±0.53 <sup>a</sup>		
BW%	21.65±2.83 <sup>cd</sup>	21.22±4.98 <sup>d</sup>	27.07±3.77 <sup>bc</sup>	23.21±0.40 <sup>cd</sup>	25.94±2.08 <sup>cd</sup>	31.52±2.27 <sup>ab</sup>	33.45±2.50 <sup>b</sup>	34.47±3.71 <sup>a</sup>	34.47±3.71 <sup>a</sup>	36.91±2.55 <sup>a</sup>		
SGR%	3.26±0.39 <sup>cd</sup>	3.20±0.70 <sup>d</sup>	3.99±0.49 <sup>bc</sup>	3.48±0.06 <sup>cd</sup>	3.84±0.28 <sup>bcd</sup>	4.56±0.28 <sup>ab</sup>	4.81±0.31 <sup>a</sup>	4.93±0.47 <sup>a</sup>	4.93±0.47 <sup>a</sup>	5.11±0.31 <sup>a</sup>		
K%	1.85±0.30	1.93±0.42	1.91±0.43	1.83±0.36	1.85±0.39	1.88±0.41	1.89±0.37	1.86±0.38	1.86±0.38	1.98±0.45		
FCR	3.14±0.39 <sup>ab</sup>	3.28±0.83 <sup>a</sup>	2.49±0.31 <sup>ede</sup>	2.91±0.05 <sup>bce</sup>	2.60±0.23 <sup>bcd</sup>	2.15±0.15 <sup>de</sup>	2.00±0.13 <sup>de</sup>	1.96±0.19 <sup>de</sup>	1.96±0.19 <sup>de</sup>	1.86±0.12 <sup>d</sup>		

\*a-d showed differences between treatments by Duncan's test.

**Table 2** Effects of Spirulina and citric acid (CA) supplementation on immunological indices of common carp in different treatments

	Control	Sp20	Sp30	CA0.5	CA1	Sp20+CA0.5	Sp20+CA1	Sp30+CA0.5	Sp30+CA1
Lysozim(u/ml/min)	30.11±2.06 <sup>a</sup>	29.33±2.08 <sup>a</sup>	27.33±2.52	30.33±2.08 <sup>b</sup>	29.00±1.00 <sup>b</sup>	28.00±2.00 <sup>b</sup>	26.00±3.46 <sup>b</sup>	32.33±3.06 <sup>b</sup>	33.00±1.73 <sup>b</sup>
Igm(mg/dl)	31.28±3.95 <sup>a</sup>	32.33±4.93 <sup>a</sup>	41.00±5.29 <sup>a</sup>	48.67±4.04 <sup>c</sup>	41.33±1.53 <sup>b</sup>	41.00±3.61 <sup>b</sup>	41.67±3.06 <sup>b</sup>	46.67±2.52 <sup>bc</sup>	40.67±1.53 <sup>b</sup>
Imonoglobulin(mg/ml)	14.06±0.27 <sup>a</sup>	15.23±0.31 <sup>ab</sup>	15.97±0.32 <sup>a</sup>	16.67±0.15 <sup>d</sup>	16.43±0.15 <sup>d</sup>	15.17±0.32 <sup>b</sup>	15.63±0.15 <sup>bc</sup>	17.20±0.30 <sup>c</sup>	16.63±0.15 <sup>d</sup>

\*a-d showed differences between treatments by Duncan's test.



The study reveals variations in the feed conversion ratios among the treatment groups. The highest recorded feed conversion ratio belongs to  $(0.48 \pm 3.28)$ , whereas the lowest level is associated with SP30+CA1  $(0.07 \pm 1.86)$ . Statistical tests indicate significant differences in feed conversion ratios among the treatments throughout the experiment ( $P < 0.05$ ). Based on the separation test, the feed conversion ratio of SP20 significantly differs from other treatments ( $P < 0.05$ ). However, it does not exhibit statistically significant differences from SP30, CA0.5, and CA1. Meanwhile, results show that SP20+CA0.5, SP20+CA1, and SP30+CA0.5 share the same feed conversion ratio ( $P > 0.05$ ).

### Immunological findings

The results of this experiment revealed that the highest lysozyme levels were observed in conjunction with Sp30+CA0.5 and Sp30+CA1, while the lowest levels were recorded for the Sp20 treatment (Table 2). A statistical analysis demonstrated a significant variance among the treatments ( $P < 0.05$ ). Additionally, it was observed that Sp30, CA0.5, CA1, Sp20+CA0.5, and Sp20+CA1 exhibited comparable lysozyme levels with no statistically significant distinctions ( $P > 0.05$ ).

According to the results, the highest IGM concentrations were found in specimens exposed to Sp30+CA0.5 and CA0.5, while the lowest concentrations were detected in those treated with Sp20 (Table 2). The statistical tests confirmed a significant distinction among the treatments ( $P < 0.05$ ). Furthermore, the findings indicated that Sp30, CA1, Sp20+CA0.5, and Sp20+CA1 exhibited equivalent IGM levels with no significant discrepancies ( $P > 0.05$ ).

The study's outcomes demonstrated that Sp30+CA0.5 displayed the highest immunoglobulin M content, whereas the lowest was associated with the Sp20 treatment. A one-way analysis of variance test revealed a statistically significant distinction between Sp30+CA0.5 and Sp20 regarding immunoglobulin M levels ( $P < 0.05$ ). Moreover, Sp30, CA0.5, CA1, Sp20+CA0.5, and Sp20+CA1 exhibited similar immunoglobulin M content (Table 2).

### Discussion

In recent years, there has been a concerted effort to replace fish meal, partially or entirely, with various plant-based sources capable of providing nutritionally dense and valuable feed. Spirulina powder, specifically *Spirulina platensis*, has emerged as a promising candidate for future aquaculture practices (Lu and Takeuchi 2004; Choonawala 2007). Due to its high protein content and essential amino acids, fish meal remains a crucial component in aquatic feeds (Güroy et al. 2022; Yousefi et al. 2022). According to the findings of this research, the growth parameters exhibited a more favorable response when higher concentrations of Spirulina and citric acid were introduced during the grow-out phase of the sample. During the initial sampling stage, Sp20+CA0.5 demonstrated the highest growth indices at the end of the eighth week. Sp30+CA0.5 and Sp30+CA1 exhibited the highest body weight index, weight gain, specific growth rate, and the lowest feed conversion ratio. Fat accumulation exceeded 15% in different concentrations, but fat levels increased with the increased inclusion of Spirulina and citric acid in Sp30+CA0.5 and Sp30+CA1. Previous research indicated that dietary lipid levels above 15% adversely affected growth and food intake in species like common carp (Jauncey 1979). In contrast, the maintenance of optimal lipid levels has been shown to yield a multitude of benefits in aquaculture, including enhanced growth rates, improved feed conversion efficiency, efficient nutrient utilization, and reduced nitrogen excretion, as demonstrated in previous studies (Yigit et al. 2002; Martin et al. 2007). The present research reinforces the significance of sustaining optimal lipid content across all treatment groups. Notably, this study highlights the promising outcomes achieved by fish groups fed with varying concentrations of Spirulina and citric acid, specifically the SP30+CA0.5 (3% Spirulina + 0.5g of citric acid per kg of diet) and SP30+CA1 (3% Spirulina + 1g of citric acid per kg of diet) treatments, which exhibited superior feed conversion efficiency and growth performance. These findings underscore the potential of nutritionally rich algae as a valuable source of fish feed, in line with the observations made by Khatoun et al. (2010). Various factors, including diet acceptability, have been recognized for their influence on the growth performance of fish species, as previously observed in catfish (*Clarius batrachus*) by Hasan et al. (1989). The current study aligns with the notion that diets incorporating algae protein can significantly enhance growth performance compared to control diets.



In the context of the present investigation, it was found that lysozyme, immunoglobulin, and Immunoglobulin M (IgM) levels exhibited notable increases, with the highest levels observed in the SP30+CA0.5 treatment group. Lysozyme activity in fish blood plays a pivotal role in assessing the innate immune system, as acknowledged by Montoya et al. (2017) and Amphan et al. (2019). Research by Reda et al. (2018) also documented increased enzyme activity in tilapia following dietary supplementation with yeast nucleotides. Furthermore, the supplementation of probiotic *Bacillus subtilis* endospores has been shown to enhance lysozyme activity in tilapias, as reported by Galagarza et al. (2018), Faheem et al. (2022), and Rosenau et al. (2022).

Numerous factors, both internal and external, can exert an influence on parameters related to the innate immune response in fish. Temperature fluctuations, stress management, and stocking density are known to have suppressive effects on these responses, while certain food additives and immunostimulants have the potential to enhance their efficiency, as documented by Magnadottir in her studies from 2006 and 2010 (Magnadottir 2006, 2010). Changes in environmental pH levels have yielded mixed results regarding immune system parameters, including lysozyme and IgM levels in the circulation. Notably, the absence of stressors in the experimental fish samples, attributed to the adequate concentrations of Spirulina and citric acid, is reflected in the heightened levels of immune-related factors.

The dietary supplementation of Spirulina and citric acid, particularly in the SP30+CA0.5 treatment, emerges as a strategy that enhances immune responses and augments disease resistance, particularly in challenging environmental conditions. Recent research corroborates these findings, demonstrating the up-regulation of immune-related genes in rainbow trout when exposed to diets containing Spirulina (Güroy et al. 2022; Yousefi et al. 2022).

## Conclusion

Based on the findings of this study, it is evident that a combination of Spirulina and citric acid can exert a more significant influence on weight gain, growth performance, and immune system enhancement. Consequently, Spirulina and citric acid can potentially enhance nutrient levels by substituting protein content. Additionally, citric acid contributes to improved absorption of vegetable protein, thereby promoting growth indices. Furthermore, incorporating Spirulina and citric acid into a fish diet can induce notable changes in various immunological factors in fish, with the most pronounced effects observed in the SP30+CA0.5 combination, leading to enhanced immune responses.

**Acknowledgments** The authors thank the individuals at the “Research and Education Center of Natural Resources and Agriculture” in Guilan Province for their cooperation in facilitating this research.

**Conflicts of interest** The authors declare no conflict of interest.

## References

- Adewolu MA (2008) Potentials of sweet potato (*Ipomoea batatas*) leaf meal as a dietary ingredient for *Tilapia zilli* fingerlings. Pak J Nutri 7:444–449
- Amphan S, Unajak S, Printrakoon C, Areechon N (2019) Feeding-regimen of  $\beta$ -glucan to enhance innate immunity and disease resistance of Nile tilapia, *Oreochromis niloticus* Linn, against *Aeromonas hydrophila* and *Flavobacterium columnare* Fish Shellfish Immunol 87:120–128
- Avron M, Ben-Amotz A (1992) Dunaliella physiology, biochemistry, and biotechnology, pp 240
- Belay A, Kato T, Ota Y (1996) Spirulina (*Arthrospira*); Potential application as an animal feed supplement. J Appl Phycol 8:303–311
- Biedenbach MA, Beuerman RW, Brown AC (1975) Graphic-digitizer analysis of axon spectra in ethmoidal and lingual branches of the trigeminal nerve. Cell Tiss Res 157:341–352
- Borowitzka MA (1997) Microalgae for aquaculture: opportunities and constraints. J Appl Phycol 9:393–401
- Brydges NM, Boulcott P, Ellis T, Braithwaite VA (2009) Quantifying stress responses induced by different handling methods in three fish species. Appl Anim Behav Sci 116:295–301
- Choonawala BB (2007) Spirulina production in brine effluent from cooling towers. Durban University of Technology, p 370
- Cross S, Debiec H, Peterlik M (1990) Mechanism and regulation of intestinal phosphate absorption. Miner Electrolyte Metab 16:115–124
- Dar BA, Khaliq R, Jha GN, Kour P, Qureshi TA (2014) Protective effects of dietary spirulina against cadmium chloride exposed histological changes in the liver of freshwater catfish *Clarias batrachus* (Linnaeus, 1758). Indian J Fish 61:83–87
- Denstadli V, Skrede A, Krogdahl A, Sahlstrom S, Storebakken T (2006) Feed intake, growth, feed conversion; digestibility, enzyme activities, and intestinal structure in Atlantic salmon (*Salmo salar* L.) fed graded levels of phytic acid. Aquaculture 256:365–376
- Dernekbası S, Unal H, Karayucel I, Aral O (2010) Effect of dietary supplementation of different rates of Spirulina (*Spirulina platensis*)



- on growth and feed conversion in the guppy (*Poecilia reticulata* Peters, 1860). *J Anim Vet Adv* 9:1395–1399
- Faheem M, Jamal R, Nazeer N, Khaliq S, Hoseinifar S, Doan H, Paolucci M (2022) Improving growth, digestive and antioxidant enzymes and immune response of juvenile grass carp (*Ctenopharyngodon idella*) by using dietary *Spirulina platensis*. *Fishes* 7:237–249
- Farag MR, Alagawany M, Abd El-Hack, ME, Dhama K (2016) Nutritional and health aspects of *Spirulina (Arthrospira)* for poultry, animals, and human. *Int J Pharmacol* 12:36–51
- Fredlund K, Isaksson M, Rossander-Hulthen L, Almgren A, Sandberg AS (2006) Absorption of zinc and calcium retention: dose-dependent inhibition by phytate. *Trace Elem Med Biol* 20:49–57
- Galagarza OA, Smith SA, Drahos DJ, Eifert JD, Williams RC, Kuhn DD (2018) Modulation of innate immunity in Nile tilapia (*Oreochromis niloticus*) by dietary supplementation of *Bacillus subtilis* endospores. *Fish Shellfish Immunol* 83:171–179
- Güroy B, Güroy D, Bilen S, Kenanoğlu ON, Şahin I, Terzi E, Karadal O, Mantoğlu S (2022) Effect of dietary *Spirulina (Arthrospira platensis)* on the growth performance, immune-related gene expression and resistance to *Vibrio anguillarum* in European seabass (*Dicentrarchus labrax*). *Aquac Res* 53:2263–2274
- Hasan MA, Alam MGM, Islam MA (1989) Evaluation of some indigenous ingredients as dietary protein sources for catfish (*Clarias batrachus*, Linn) fry. In E.A. Huisman, K. Conneveld & A.H.M Bouwmans (Ed.), *Aquac Res in Asia: Management techniques and nutrition*, Wageningen, Pudoc, pp 125–137
- Higgs DA, Dosañh BS, Prendergast AF, Beames RM, Hardy RW, Riley W, Deacon G (1995) Use of rapeseed/canola protein products in finfish diets. In CE Lim and DJ Sessa (Eds.), *Nutrition and Utilization Technology in Aquaculture*, pp130–156
- Jackson AJ, Capper BS, Matty AJ (1982) Evaluation of some plant proteins in complete diets for the tilapia (*Sarotherodon mossambicus*). *Aquaculture* 27:97–109
- James R, Sampath K, Nagarajan R, Vellaisamy P, Manikandan MM (2009) Effect of dietary spirulina on reduction of copper toxicity and improvement of growth, blood parameters and phosphatases activities in carp, *Cirrhinus mrigala* (Hamilton 1822). *Indian J Exp Biol* 47:754–759
- Jauncey K (1979) Growth and nutrition of carp in heated effluents. University of Ashton in Birmingham, Birmingham. PhD thesis, P 202
- Kaushik SJ (1990) Use alternative protein sources for intensively rearing carnivorous fish. In R Flos, L Tort and P Torres (Ed.), *Mediterranean Aquaculture*, pp125–138
- Khatoun N, Chattopadhyay P, Mukhopadhyay A, Mukhopadhyay M, Pal R (2009) Algae diet in Prawn Aquaculture. *Fish Chimes* 28:44–47
- Khatoun N, Chaudhuri A, Sen Roy S, Kundu N, Mukherjee S, Majumdar D, Homechaudhuri S, Pal R, (2010) Algae as a feed supplement in fish nutrition. *J Botl Soc* 64:85–93
- Kim SS, Shin SJ, Han HS, Kim JD, Lee KJ (2015) Effects of dietary *Spirulina pacifica* on innate immunity and disease resistance against *Edwardsiella tarda* in olive flounder *Paralichthys olivaceus*. *Isr J Aquac* 67:1–9
- Kumar A, Bhatnagar A, Garg SK, Jana SN (2010) Growth performance of Nile tilapia, *Oreochromis niloticus* (Linn.) about the provision of substrate and supplementary feeding and grown in brackish water ponds. *Asian Fish Sci* 22:1211–1233
- Lee YK (1997) Commercial microalgae production in the Asia Pacific rim. *J Appl Phycol* 9:403–411
- Lu J, Takeuchi T (2004) Spawning and egg quality of the tilapia *Oreochromis niloticus* fed solely on raw *Spirulina* throughout three generations. *Aquaculture* 234:625–640
- Magnadottir B (2006) Innate immunity of fish (overview). *Fish Shellfish Immunol* 120:137–151
- Magnadottir B (2010) Immunological control of fish diseases. *Mar Biotechnol* 12:361–379
- Martins DA, Valente LMP, Lall SP (2007) Effects of dietary lipid level on growth and lipid utilization by juvenile Atlantic halibut (*Hippoglossus hippoglossus*, L). *Aquaculture* 263:150–158
- Montoya LNF, Martins TP, Gimbo RY, Zanuzzo FS, Urbinati EC (2017)  $\beta$ -Glucan-induced cortisol levels improve the early immune response in matrinxã (Brycon amazonicus). *Fish Shellfish Immunol* 60:197–204
- Moreno-Garcia L, Adjall'e K, Barnab'e S, Raghavan, GSV (2017) Microalgae biomass production for a biorefinery system: recent advances and the way towards sustainability. *Renew Sust Energ Rev* 76:493–506
- Navarro N, Sarasquete C (1998) Use of freeze-dried microalgae for rearing gilthead seabream, *Sparus aurata*, larvae. Growth, histology, and water quality. *Aquaculture* 167:179–93
- Priyadarshani I, Rath B (2012) Commercial and industrial applications of microalgae a review. *J Algal Biomass* 3:89–100
- Rahimnejad S (2013) Partial replacement of fish meal with spirulina pacifica in diets for parrot fish (*Oplegnathus fasciatus*). *Turk J Fish Aquat Sci* 13:197–204
- Reda RM, Selim KM, Mahmoud R, El-Araby IE (2018) Effect of dietary yeast nucleotide on antioxidant activity, non-specific immunity, intestinal cytokines, and disease resistance in Nile Tilapia. *Fish Shellfish Immunol* 80:281–290
- Rosenau S, Ciulu M, Reimer C, Mott A, Tetens J, Mörlein D (2022) Feeding green: *Spirulina (Arthrospira platensis)* induced changes in production performance and quality of salmonid species. *Aquac Res* 53:4276–4287
- Roy SS, Pal R (2015) Microalgae in aquaculture: a review with particular references to nutritional value and fish dietetics. *Proceed Zool Soc* 68:1–8
- Sarker MSA, Satoh S, Kiron V (2005) Supplementation of citric acid and amino acid-chelated trace element to develop environment-friendly feed for red sea bream, *Pagrus major*. *Aquaculture* 248:3–11
- Sarker PK, Gamble MM, Kelson S, Kapuscinski AR (2016) Nile tilapia (*Oreochromis niloticus*) show high digestibility of lipid and fatty acids from marine *Schizochytrium sp.*, and of protein and essential amino acids from freshwater *Spirulina sp.* Feed ingredients. *Aquac Nutr* 22:109–119
- Storebakken T, Shearer KD, Roem AJ (1998) Availability of protein, phosphorus and other elements in fish meal, soy protein concentrate, and phytase treated soy-protein-concentrate based diets to Atlantic salmon, *Salmo salar*. *Aquaculture* 161:365–379
- Sugiura SH, Gabaudan J, Dong FM, Hardy RW (2001) Dietary microbial phytase supplementation and the utilization of phosphorus, trace minerals and protein by rainbow trout (*Oncorhynchus mykiss*, W) fed soybean-meal based diets. *Aquac Res* 32:583–592
- Venkataraman LV (1997) *Spirulina platensis (Arthrospira)*: Physiology, cell biology and biotechnology. *J Appl Phycol* 9:295–296
- Vielma J, Ruohonen K, Lall SP (1999) Supplemental citric acid and particle size of fish bone-meal influence the availability of min-





- erals in rainbow trout (*Oncorhynchus mykiss*, W). *Aquac Nutr* 5:65–71
- Wood RJ, Serfaty-Lacrosniere C (1992) Gastric acidity, atrophic gastritis, and calcium absorption. *Nutr Rev* 50:33–40
- Yamaguchi K (1997) Recent advances in microalgal bioscience in Japan, with particular reference to utilization of biomass and metabolites: A review. *J Appl Phycol* 8:227–233
- Yigit M, Yardim O, Koshio S (2002) The protein sparing effects of high lipid levels in diets for rainbow trout (*Oncorhynchus mykiss*, W) with particular reference to reduction of total nitrogen excretion. *Isr J Aquac Bamidgeh* 54:79–88
- Yousefi M, Ahmadifar M, Mohammadzadeh S, Kalhor N, EslimiEsfahani D, Bagheri A, Mashhadizadeh N, ShahriariMoghadam M, Ahmadifar E (2022) Individual and combined effects of the dietary *Spirulina platensis* and *Bacillus licheniformis* supplementation on growth performance, antioxidant capacity, innate immunity, relative gene expression and resistance of goldfish, *Carassius auratus* to *Aeromonas hydrophila*. *Fish Shellfish Immunol* 27:1070–1078
- Zyla K, Ledoux DR, Garcia A, Veum TL (1995) An in vitro procedure for studying enzymic dephosphorylation of phytate in maize–soybean feeds for Turkey poults. *Br J Nutr* 74:3–17

**Publisher's Note**

IAU remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

